

## Product datasheet for TA802703M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Progesterone Receptor (PGR) Mouse Monoclonal Antibody [Clone ID: OTI15C1]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI15C1

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

**Isotype:** IgG2b

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-298 of human PGR

(NP\_000917) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 98.8 kDa

**Gene Name:** progesterone receptor

Database Link: NP 000917

Entrez Gene 5241 Human

P06401





**Background:** This gene encodes a member of the steroid receptor superfamily. The encoded protein

mediates the physiological effects of progesterone, which plays a central role in reproductive events associated with the establishment and maintenance of pregnancy. This gene uses two distinct promotors and translation start sites in the first exon to produce two isoforms, A and B. The two isoforms are identical except for the additional 165 amino acids found in the N-terminus of isoform B and mediate their own response genes and physiologic effects with


little overlap. [provided by RefSeq, Jan 2011]

Synonyms: NR3C3; PR

**Protein Families:** Druggable Genome, Nuclear Hormone Receptor, Transcription Factors

**Protein Pathways:** Oocyte meiosis, Progesterone-mediated oocyte maturation

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PGR ([RC221303], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PGR. (1:5. Positive lysates [LY424456] (100ug) and [LC424456] (20ug) can be purchased separately from OriGene.