

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA802710S

Progesterone Receptor (PGR) Mouse Monoclonal Antibody [Clone ID: OTI7A9]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI7A9	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Human recombinant protein fragment corresponding to amino acids 1-298 of human PGR (NP_000917) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	98.8 kDa	
Gene Name:	progesterone receptor	
Database Link:	<u>NP_000917</u> <u>Entrez Gene 5241 Human</u> <u>P06401</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	Progesterone Receptor (PGR) Mouse Monoclonal Antibody [Clone ID: OTI7A9] – TA802710S	
Background:	This gene encodes a member of the steroid receptor superfamily. The encoded protein mediates the physiological effects of progesterone, which plays a central role in reproductive events associated with the establishment and maintenance of pregnancy. This gene uses two distinct promotors and translation start sites in the first exon to produce two isoforms, A and B. The two isoforms are identical except for the additional 165 amino acids found in the N- terminus of isoform B and mediate their own response genes and physiologic effects with little overlap. [provided by RefSeq, Jan 2011]	
Synonyms:	NR3C3; PR	
Protein Families:	Druggable Genome, Nuclear Hormone Receptor, Transcription Factors	
Protein Pathway	S: Oocyte meiosis, Progesterone-mediated oocyte maturation	

Product images:

%// _

170 —	100
	-
130 —	
100 —	
	Contraction of the local division of the loc
70 —	
55 -	
	-
40 -	
35	
25 -	
15 —	
10 -	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PGR ([RC221303], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PGR. (1:5. Positive lysates [LY424456] (100ug) and [LC424456] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US