

Product datasheet for TA802776BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

beta 2 Microglobulin (B2M) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI1A9]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A9

Applications: ELISA, IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Purified native Human beta 2 Microglobulin protein

Formulation: PBS (pH 7.4) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 11.7 kDa

Gene Name: beta-2-microglobulin

Database Link: NP 004039

Entrez Gene 567 Human

P61769

Background: This gene encodes a serum protein found in association with the major histocompatibility

complex (MHC) class I heavy chain on the surface of nearly all nucleated cells. The protein has

a predominantly beta-pleated sheet structure that can form amyloid fibrils in some

pathological conditions. A mutation in this gene has been shown to result in hypercatabolic

hypoproteinemia. [provided by RefSeq, Sep 2009]

Synonyms: IMD43






Protein Families: Druggable Genome, Secreted Protein


Protein Pathways: Antigen processing and presentation

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY B2M ([RC207587], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-B2M. Positive lysates [LY401311] (100ug) and [LC401311] (20ug) can be purchased separately from OriGene.


Beta 2-Microglobulin



Standard curve for ELISA analysis with B2M recombinant protein (dilution range from 0.3ng/ml to 200ng/ml) using B2M Capture Antibody (Cat# [TA803600]/[TA803617]/[TA802776]) at 5ug/ml and HRP conjugated B2M Detection mAb (Cat# [TA590171]) at 0.125ug/ml.







Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-B2M mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802776])

Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-B2M mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802776])