

## **Product datasheet for TA802840**

### 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# CD63 Mouse Monoclonal Antibody [Clone ID: OTI5D9]

**Product data:** 

**Product Type:** Primary Antibodies

Clone Name: OTI5D9
Applications: FC, WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CD63 (NP\_001771) produced in HEK293T

cell

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 25.5 kDa

Gene Name: CD63 molecule

Database Link: NP 001771

Entrez Gene 967 Human

P08962



#### Background:


The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]

Synonyms: LAMP-3; ME491; MLA1; OMA81H; TSPAN30


**Protein Families:** Druggable Genome, Transmembrane

**Protein Pathways:** Lysosome

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD63 ([RC201733], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD63. Positive lysates [LY419757] (100ug) and [LC419757] (20ug) can be purchased separately from OriGene.




Western blot analysis of extracts (35ug) from MCF7 cell line by using anti-CD63 monoclonal antibody (1:500).






HEK293T cells transfected with either [RC201733] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CD63 antibody (TA802840), and then analyzed by flow cytometry (1:100).



Flow cytometric Analysis of living A549 cells, using anti-CD63 antibody (TA802840), (Red), compared to a nonspecific negative control antibody, (Blue).



Flow cytometric Analysis of MCF-7 cells, using anti-CD63 antibody (TA802840), (Red), compared to isotype control, (green), and negative control (PBS), (Blue) (1:100)