

Product datasheet for TA802929S

DAP3 Mouse Monoclonal Antibody [Clone ID: OTI3H6]

Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3H6
Applications:	WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 161-398 of human DAP3 (NP_387506) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	45.4 kDa
Gene Name:	death associated protein 3
Database Link:	<u>NP 387506</u>
	<u>Entrez Gene 7818 Human</u> <u>P51398</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Serigene DAP3 Mouse Monoclonal Antibody [Clone ID: OTI3H6] – TA802929S

Background:Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in
protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes)
consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein
to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed.
Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that
the latter contain a 5S rRNA. Among different species, the proteins comprising the
mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which
prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein
that also participates in apoptotic pathways which are initiated by tumor necrosis factor-
alpha, Fas ligand, and gamma interferon. This protein S27. Multiple alternatively spliced
transcript variants encoding distinct isoforms have been found for this gene. Pseudogenes
corresponding to this gene are found on chromosomes 1q and 2q. [provided by RefSeq, Dec
2010]

Synonyms:bMRP-10; DAP-3; MRP-S29; MRPS29Protein Families:Druggable Genome

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY DAP3 ([RC223182], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DAP3. Positive lysates [LY403256] (100ug) and [LC403256] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US