

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA802955BM

## CD68 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8A4]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI8A4
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human
Host:	Mouse
lsotype:	lgG2b
Clonality:	Monoclonal
Immunogen:	Human recombinant protein fragment corresponding to amino acids 22-319 of human CD68 (NP_001242)produced in SF9 cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	37.2 kDa
Gene Name:	CD68 molecule
Database Link:	<u>NP_001242</u> <u>Entrez Gene 968 Human</u> <u>P34810</u>



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

#### CD68 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8A4] – TA802955BM CD68 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI8A4] – TA802955BM

Background:This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human<br/>monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated<br/>membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and<br/>endosomes with a smaller fraction circulating to the cell surface. It is a type I integral<br/>membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and<br/>organ-specific lectins or selectins. The protein is also a member of the scavenger receptor<br/>family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis,<br/>and mediate the recruitment and activation of macrophages. Alternative splicing results in<br/>multiple transcripts encoding different isoforms. [provided by RefSeq, Jul 2008]


Synonyms: GP110; LAMP4; SCARD1

Lysosome


Protein Families:

Protein Pathways:

### **Product images:**




Druggable Genome, Transmembrane



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD68 ([RC200392], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD68. Positive lysates [LY420047] (100ug) and [LC420047] (20ug) can be purchased separately from OriGene.

Equivalent amounts of cell lysates (10 ug per lane) of wild-type Hela cells (WT, Cat# LC810HELA) and CD68-Knockout Hela cells (KO, Cat# [LC810037]) were separated by SDS-PAGE and immunoblotted with anti-CD68 monoclonal antibody [TA802955]. Then the blotted membrane was stripped and reprobed with antib-actin antibody ([TA811000]) as a loading control (1:500).


This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US 

Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-CD68 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

Immunohistochemical staining of paraffinembedded Carcinoma of Human lung tissue using anti-CD68 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-CD68 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-CD68 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-CD68 mouse monoclonal antibody. (Heatinduced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-CD68 mouse monoclonal antibody. (Heat-induced epitope retrieval by 10mM citric buffer, pH6.0, 120°C for 3min, [TA802955])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US