

# Product datasheet for TA802983S

# CD68 Mouse Monoclonal Antibody [Clone ID: OTI4D8]

# **Product data:**

#### **Product Type: Primary Antibodies Clone Name:** OTI4D8 **Applications:** WB Recommended Dilution: WB 1:2000 **Reactivity:** Human Host: Mouse Isotype: lgG2a **Clonality:** Monoclonal Human recombinant protein fragment corresponding to amino acids 22-319 of human CD68 Immunogen: (NP 001242)produced in SF9 cell. Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. **Concentration:** 1 mg/ml **Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) **Conjugation:** Unconjugated Store at -20°C as received. Storage: Stability: Stable for 12 months from date of receipt. Gene Name: CD68 molecule Database Link: NP 001242 Entrez Gene 968 Human P34810



View online »

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn


### **GRIGENE** CD68 Mouse Monoclonal Antibody [Clone ID: OTI4D8] – TA802983S

Background: This gene encodes a 110-kD transmembrane glycoprotein that is highly expressed by human monocytes and tissue macrophages. It is a member of the lysosomal/endosomal-associated membrane glycoprotein (LAMP) family. The protein primarily localizes to lysosomes and endosomes with a smaller fraction circulating to the cell surface. It is a type I integral membrane protein with a heavily glycosylated extracellular domain and binds to tissue- and organ-specific lectins or selectins. The protein is also a member of the scavenger receptor family. Scavenger receptors typically function to clear cellular debris, promote phagocytosis, and mediate the recruitment and activation of macrophages. Alternative splicing results in multiple transcripts encoding different isoforms. [provided by RefSeq, Jul 2008]

Synonyms: GP110; LAMP4; SCARD1

Protein Families:Druggable Genome, TransmembraneProtein Pathways:Lysosome

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD68 ([RC200392], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD68. Positive lysates [LY420047] (100ug) and [LC420047] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US