

Product datasheet for TA803095M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RIP3 (RIPK3) Mouse Monoclonal Antibody [Clone ID: OTI2F11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2F11

Applications: WB

Recommended Dilution: WB 1:200 - 1:1000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 233-518 of human

RIPK3 (NP_006862) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 56.7 kDa

Gene Name: receptor interacting serine/threonine kinase 3

Database Link: NP 006862

Entrez Gene 11035 Human

O9Y572






Background:

The product of this gene is a member of the receptor-interacting protein (RIP) family of serine/threonine protein kinases, and contains a C-terminal domain unique from other RIP family members. The encoded protein is predominantly localized to the cytoplasm, and can undergo nucleocytoplasmic shuttling dependent on novel nuclear localization and export signals. It is a component of the tumor necrosis factor (TNF) receptor-I signaling complex, and can induce apoptosis and weakly activate the NF-kappaB transcription factor. [provided by RefSeq, Jul 2008]


Synonyms: RIP3

Protein Families: Druggable Genome, Protein Kinase
Protein Pathways: Cytosolic DNA-sensing pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RIPK3 ([RC209549], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RIPK3.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-RIPK3 monoclonal antibody at 1:200.