

Product datasheet for TA803267M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytokeratin 14 (KRT14) Mouse Monoclonal Antibody [Clone ID: OTI4A7]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A7
Applications: IHC, WB
Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 261-472 of human

KRT14 (NP_000517) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 51.4 kDa

Gene Name: keratin 14

Database Link: NP 000517

Entrez Gene 16664 MouseEntrez Gene 287701 RatEntrez Gene 3861 Human

P02533

Background: This gene encodes a member of the keratin family, the most diverse group of intermediate


filaments. This gene product, a type I keratin, is usually found as a heterotetramer with two keratin 5 molecules, a type II keratin. Together they form the cytoskeleton of epithelial cells. Mutations in the genes for these keratins are associated with epidermolysis bullosa simplex. At least one pseudogene has been identified at 17p12-p11. [provided by RefSeq, Jul 2008]






Synonyms: CK14; EBS3; EBS4; K14; NFJ

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KRT14 ([RC214907], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KRT14.




Western blot analysis of extracts (35ug) from A431 cells by using anti-KRT14 monoclonal antibody (1:500).






Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-KRT14 mouse monoclonal antibody. ([TA803267]) Dilution: 1:150. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-KRT14 mouse monoclonal antibody. ([TA803267]) Dilution: 1:150. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Carcinoma of Human bladder tissue using anti-KRT14 mouse monoclonal antibody. ([TA803267]) Dilution: 1:150. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.