

## Product datasheet for **TA803393M**

### CD63 Mouse Monoclonal Antibody [Clone ID: OTI3D9]

#### Product data:

Product Type:	Primary Antibodies
Clone Name:	OTI3D9
Applications:	FC, WB
Recommended Dilution:	WB 1:2000
Reactivity:	Human
Host:	Mouse
Isotype:	IgG2b
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human CD63 (NP_001771) produced in HEK293T cell.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.
Concentration:	1 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25.5 kDa
Gene Name:	CD63 molecule
Database Link:	<a href="#">NP_001771</a> <a href="#">Entrez Gene 967 Human</a> <a href="#">P08962</a>



[View online »](#)

**Background:**

The protein encoded by this gene is a member of the transmembrane 4 superfamily, also known as the tetraspanin family. Most of these members are cell-surface proteins that are characterized by the presence of four hydrophobic domains. The proteins mediate signal transduction events that play a role in the regulation of cell development, activation, growth and motility. The encoded protein is a cell surface glycoprotein that is known to complex with integrins. It may function as a blood platelet activation marker. Deficiency of this protein is associated with Hermansky-Pudlak syndrome. Also this gene has been associated with tumor progression. Alternative splicing results in multiple transcript variants encoding different protein isoforms. [provided by RefSeq, Apr 2012]

**Synonyms:**


LAMP-3; ME491; MLA1; OMA81H; TSPAN30

**Protein Families:**


Druggable Genome, Transmembrane

**Protein Pathways:**


Lysosome

**Product images:**


HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CD63 ([RC201733], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD63. Positive lysates [LY419757] (100ug) and [LC419757] (20ug) can be purchased separately from OriGene.



HEK293T cells transfected with either [RC201733] overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-CD63 antibody ([TA803393]), and then analyzed by flow cytometry (1:100).



Flow cytometric Analysis of MCF-7 cells, using anti-CD63 antibody ([TA803393]), (Red), compared to isotype control, (green), and negative control (PBS), (Blue) (1:100)