

### Product datasheet for TA803407AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## MAP1LC3A Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI2D5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human MAP1LC3A (NP\_115903) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 14.1 kDa

**Gene Name:** microtubule associated protein 1 light chain 3 alpha

Database Link: NP 115903

Entrez Gene 66734 MouseEntrez Gene 362245 RatEntrez Gene 84557 Human

Q9H492

**Background:** MAP1A and MAP1B are microtubule-associated proteins which mediate the physical

interactions between microtubules and components of the cytoskeleton. MAP1A and MAP1B each consist of a heavy chain subunit and multiple light chain subunits. The protein encoded by this gene is one of the light chain subunits and can associate with either MAP1A or MAP1B.

Two transcript variants encoding different isoforms have been found for this gene. The expression of variant 1 is suppressed in many tumor cell lines, suggesting that may be

involved in carcinogenesis. [provided by RefSeq, Feb 2012]






Synonyms:

ATG8E; LC3; LC3A; MAP1ALC3; MAP1BLC3

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MAP1LC3A ([RC202222], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MAP1LC3A.