

## Product datasheet for TA803552AM

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### MRPS7 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI5C8]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5C8

**Applications:** WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 38-242 of human

MRPS7 (NP\_057055) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Predicted Protein Size: 28 kDa

**Gene Name:** mitochondrial ribosomal protein S7

Database Link: NP 057055

Entrez Gene 51081 Human

O9Y2R9






### Background:

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 28S subunit protein. In the prokaryotic ribosome, the comparable protein is thought to play an essential role in organizing the 3' domain of the 16 S rRNA in the vicinity of the P- and A-sites. Pseudogenes corresponding to this gene are found on chromosomes 8p and 12p. [provided by RefSeq, Jul


Synonyms:

bMRP27a; MRP-S; MRP-S7; RP-S7; RPMS7; S7mt

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY MRPS7 ([RC200031], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-MRPS7. Positive lysates [LY414276] (100ug) and [LC414276] (20ug) can be purchased separately from OriGene.



Equivalent amounts of cell lysates (10 ug per lane) of wild-type HeLa cells (WT, Cat# LC810HELA) and MRPS7-Knockout HeLa cells (KO, Cat# [LC831777]) were separated by SDS-PAGE and immunoblotted with anti-MRPS7 monoclonal antibody [TA803552] (1:250). Then the blotted membrane was stripped and reprobed with anti-HSP90 antibody as a loading control.