

Product datasheet for TA803648

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200 Rockville. MD 20850. US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

IRAK2 Mouse Monoclonal Antibody [Clone ID: OTI4C1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4C1
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 362-625 of human

IRAK2 (NP_001561) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: interleukin 1 receptor associated kinase 2

Database Link: NP 001561

Entrez Gene 3656 Human

O43187

Background: IRAK2 encodes the interleukin-1 receptor-associated kinase 2, one of two putative

serine/threonine kinases that become associated with the interleukin-1 receptor (IL1R) upon stimulation. IRAK2 is reported to participate in the IL1-induced upregulation of NF-kappaB.

[provided by RefSeq, Jul 2008]

Synonyms: IRAK-2


Protein Families: Druggable Genome, Protein Kinase






Protein Pathways: Apoptosis, Neurotrophin signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY IRAK2 (Cat# [RC218437], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IRAK2(Cat# TA803648).



Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human breast tissue using anti-IRAK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, TA803648)