

Product datasheet for TA803720S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TACC2 Mouse Monoclonal Antibody [Clone ID: OTI3G1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3G1
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 727-1026 of human

TACC2 (NP_008928) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: transforming acidic coiled-coil containing protein 2

Database Link: NP 008928

Entrez Gene 10579 Human

<u>095359</u>

Background: Transforming acidic coiled-coil proteins are a conserved family of centrosome- and

microtubule-interacting proteins that are implicated in cancer. This gene encodes a protein

that concentrates at centrosomes throughout the cell cycle. This gene lies within a

chromosomal region associated with tumorigenesis. Expression of this gene is induced by erythropoietin and is thought to affect the progression of breast tumors. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jul


2008]






Synonyms: AZU-1; ECTACC

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY TACC2 (Cat# [RC215999], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TACC2(Cat# [TA803720]).



Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-TACC2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA803720])