

Product datasheet for TA803726M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TACC2 Mouse Monoclonal Antibody [Clone ID: OTI1C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C3
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 727-1026 of human

TACC2 (NP_008928) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: transforming acidic coiled-coil containing protein 2

Database Link: NP 008928


Entrez Gene 10579 Human

<u>095359</u>


Synonyms: AZU-1; ECTACC



Product images:




Human recombinant protein fragment corresponding to amino acids 727-1026 of human TACC2 (NP_008928) produced in E.coli.




Immunohistochemical staining of paraffinembedded Human skin tissue using anti-TACC2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Human adult heart tissue using anti-TACC2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human embryonic liver tissue using anti-TACC2 mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human placenta tissue using anti-TACC2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.