

Product datasheet for TA803992BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Thyroglobulin (TG) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI11F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI11F8

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 20-358 of human TG

(NP_003226) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: thyroglobulin

Database Link: NP 003226

Entrez Gene 7038 Human

P01266





Thyroglobulin (TG) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI11F8] – TA803992BM

Background:


Thyroglobulin (Tg) is a glycoprotein homodimer produced predominantly by the thryroid gland. It acts as a substrate for the synthesis of thyroxine and triiodothyronine as well as the storage of the inactive forms of thyroid hormone and iodine. Thyroglobulin is secreted from the endoplasmic reticulum to its site of iodination, and subsequent thyroxine biosynthesis, in the follicular lumen. Mutations in this gene cause thyroid dyshormonogenesis, manifested as goiter, and are associated with moderate to severe congenital hypothyroidism. Polymorphisms in this gene are associated with susceptibility to autoimmune thyroid diseases (AITD) such as Graves disease and Hashimoto thryoiditis. [provided by RefSeq, Nov 2009]

Synonyms: AITD3; TGN

Protein Families: Druggable Genome

Protein Pathways: Autoimmune thyroid disease

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY TG ([RC216216], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-TG.