

## Product datasheet for TA804152M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## **ACE2 Mouse Monoclonal Antibody [Clone ID: OTI1G4] (Angiotensin Converting Enzyme 2)**

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1G4
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 18-237 of human ACE2

(NP\_068576) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** angiotensin I converting enzyme 2

Database Link: NP 068576

Entrez Gene 59272 Human

Q9BYF1



# ACE2 Mouse Monoclonal Antibody [Clone ID: OTI1G4] (Angiotensin Converting Enzyme 2) – TA804152M

Background:


The protein encoded by this gene belongs to the angiotensin-converting enzyme family of dipeptidyl carboxydipeptidases and has considerable homology to human angiotensin 1 converting enzyme. This secreted protein catalyzes the cleavage of angiotensin I into angiotensin 1-9, and angiotensin II into the vasodilator angiotensin 1-7. The organ- and cell-specific expression of this gene suggests that it may play a role in the regulation of cardiovascular and renal function, as well as fertility. In addition, the encoded protein is a functional receptor for the spike glycoprotein of the human coronaviruses SARS and HCoV-NL63. [provided by RefSeq, Jul 2008]

Synonyms: ACEH

**Protein Families:** Druggable Genome, Secreted Protein, Transmembrane

**Protein Pathways:** Renin-angiotensin system

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ACE2 ([RC208442], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ACE2 ([TA804152]). (1:500)




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.






Immunohistochemical staining of paraffinembedded Human Kidney tissue using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heatinduced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120°C for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded Human renal cell carcinoma tissue using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.




Immunohistochemical staining of paraffinembedded carcinoma of Human lung tissue using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human testicular tissue within the normal limits using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human Gastric Carcinoma using anti-ACE2 mouse monoclonal antibody ([TA804152]). Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.0) at 120 oC for 3 min. (1:800). Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.