

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA804388M

Centrin 1 (CETN1) Mouse Monoclonal Antibody [Clone ID: OTI7G7]

Product data:

| Product Type: | Primary Antibodies |
|--------------------------|---|
| Clone Name: | OTI7G7 |
| Applications: | WB |
| Recommended Dilution: | WB 1:2000 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Mouse |
| lsotype: | lgG1 |
| Clonality: | Monoclonal |
| Immunogen: | Full length human recombinant protein of human CETN1 (NP_004057) produced in E.coli. |
| Formulation: | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide. |
| Concentration: | 1 mg/ml |
| Purification: | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G) |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C as received. |
| Stability: | Stable for 12 months from date of receipt. |
| Predicted Protein Size: | 19.4 kDa |
| Gene Name: | centrin 1 |
| Database Link: | <u>NP_004057</u> <u>Entrez Gene 1068 Human</u> <u>Q12798</u> |
| Background: Synonyms: | The protein encoded by this gene plays important roles in the determination of centrosome position and segregation, and in the process of microtubule severing. This encoded protein is localized to the centrosome of interphase cells, and redistributes to the region of the spindle poles during mitosis, reflecting the dynamic behavior of the centrosome during the cell cycle. [provided by RefSeq, Jul 2008] CEN1; CETN |
| Synonymis. | |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Protein Families: Druggable Genome

Product images:

| 170 | _ | |
|-----|---|---|
| 130 | _ | |
| 100 | _ | |
| 70 | _ | |
| 55 | | |
| 40 | | |
| 35 | — | |
| 25 | — | |
| 15 | — | - |
| 10 | — | |

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CETN1 ([RC206285], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CETN1. Positive lysates [LY401316] (100ug) and [LC401316] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US