

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA804470BM

## PAFAH1B3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4A9]

## **Product data:**

Product Type:	Primary Antibodies
Clone Name:	OTI4A9
Applications:	IHC, WB
Recommended Dilution:	WB 1:2000, IHC 1:150
Reactivity:	Human, Mouse, Rat
Host:	Mouse
lsotype:	lgG1
Clonality:	Monoclonal
Immunogen:	Full length human recombinant protein of human PAFAH1B3 (NP_001139411) produced in E.coli.
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol.
Concentration:	0.5 mg/ml
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)
Conjugation:	HRP
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Predicted Protein Size:	25.6 kDa
Gene Name:	platelet activating factor acetylhydrolase 1b catalytic subunit 3
Database Link:	<u>NP_001139411</u> <u>Entrez Gene 5050 Human</u> <u>Q15102</u>




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### Serigene PAFAH1B3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI4A9] – TA804470BM


Background:This gene encodes an acetylhydrolase that catalyzes the removal of an acetyl group from the<br/>glycerol backbone of platelet-activating factor. The encoded enzyme is a subunit of the<br/>platelet-activating factor acetylhydrolase isoform 1B complex, which consists of the catalytic<br/>beta and gamma subunits and the regulatory alpha subunit. This complex functions in brain<br/>development. A translocation between this gene on chromosome 19 and the CDC-like kinase<br/>2 gene on chromosome 1 has been observed, and was associated with mental retardation,<br/>ataxia, and atrophy of the brain. Alternatively spliced transcript variants have been described.<br/>[provided by RefSeq, Mar 2009]

Synonyms:	PAFAHG
Protein Families:	Druggable Genome
Protein Pathways:	Ether lipid metabolism, Metabolic pathways

### **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PAFAH1B3 ([RC227227], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PAFAH1B3. Positive lysates [LY429044] (100ug) and [LC429044] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human kidney tissue using anti-PAFAH1B3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA804470])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US



Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-PAFAH1B3 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA804470])

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US