

Product datasheet for TA804515

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

DCC Mouse Monoclonal Antibody [Clone ID: OTI4A1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A1
Applications: WB

Recommended Dilution: WB 1:200

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human DCC (NP_005206) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 158.3 kDa

Gene Name: DCC netrin 1 receptor

Database Link: NP 005206

Entrez Gene 13176 MouseEntrez Gene 25311 RatEntrez Gene 1630 Human

P43146





Background:


This gene encodes a netrin 1 receptor. The transmembrane protein is a member of the immunoglobulin superfamily of cell adhesion molecules, and mediates axon guidance of neuronal growth cones towards sources of netrin 1 ligand. The cytoplasmic tail interacts with the tyrosine kinases Src and focal adhesion kinase (FAK, also known as PTK2) to mediate axon attraction. The protein partially localizes to lipid rafts, and induces apoptosis in the absence of ligand. The protein functions as a tumor suppressor, and is frequently mutated or downregulated in colorectal cancer and esophageal carcinoma. [provided by RefSeq, Oct 2009]

Synonyms: CRC18; CRCR1; IGDCC1; MRMV1; NTN1R1

Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Axon guidance, Colorectal cancer, Pathways in cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY DCC (Cat# [RC218506], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-DCC(Cat# TA804515). Positive lysates [LY417439] (100ug) and [LC417439] (20ug) can be purchased separately from OriGene.