

Product datasheet for TA804802M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

SNX12 Mouse Monoclonal Antibody [Clone ID: OTI8A11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8A11
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 **Reactivity:** Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human SNX12 (NP_037478) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 18.7 kDa

Gene Name: sorting nexin 12

Database Link: NP 037478

Entrez Gene 55988 MouseEntrez Gene 363478 RatEntrez Gene 29934 Human

Q9UMY4

Background: This gene encodes a member of the sorting nexin family. Members of this family contain a

phox (PX) domain, which is a phosphoinositide binding domain, and are involved in

intracellular trafficking. This protein does not contain a coiled coil region, like some family members. A similar protein in mouse may be involved in regulating the neurite outgrowth.


Alternate splicing results in multiple transcript variants. [provided by RefSeq, Jan

Synonyms: MGC118982; MGC118983






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY SNX12 ([RC224303], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-SNX12. Positive lysates [LY415650] (100ug) and [LC415650] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-SNX12 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.