

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA805008AM

## HDAC4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1E12]

#### **Product data:**

Product Type:	Primary Antibodies	
Clone Name:	OTI1E12	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG1	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human HDAC4 (NP_006028) produced in HEK293T cell.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	0.5 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Biotin	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	118.9 kDa	
Gene Name:	histone deacetylase 4	
Database Link:	<u>NP_006028</u> <u>Entrez Gene 208727 MouseEntrez Gene 363287 RatEntrez Gene 9759 Human</u> <u>P56524</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

	HDAC4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1E12] – TA805008AM
Background:	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. The protein encoded by this gene belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3. [provided by RefSeq, Jul 2008]
Synonyms: Protein Families	AHO3; BDMR; HA6116; HD4; HDAC-4; HDAC-A; HDACA Druggable Genome, Transcription Factors

### **Product images:**

170	-	
130	-	
100		
70	-	-
55		
40		
35	-	
25	-	
15	-	
10	-1	

HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY HDAC4 (Cat# [RC211495], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-HDAC4 (Cat# [TA805008]). Positive lysates [LY416910] (100ug) and [LC416910] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US