

# **Product datasheet for TA805047M**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# CD56 (NCAM1) Mouse Monoclonal Antibody [Clone ID: OTI7D11]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7D11

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 20-718 of human

NCAM1 (NP\_851996) produced in HEK293T cell.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 94.4 kDa

**Gene Name:** neural cell adhesion molecule 1

Database Link: NP 851996

Entrez Gene 17967 MouseEntrez Gene 24586 RatEntrez Gene 4684 Human

P13591

**Background:** This gene encodes a cell adhesion protein which is a member of the immunoglobulin

superfamily. The encoded protein is involved in cell-to-cell interactions as well as cell-matrix interactions during development and differentiation. The encoded protein has been shown to be involved in development of the nervous system, and for cells involved in the expansion of T cells and dendritic cells which play an important role in immune surveillance. Alternative

splicing results in multiple transcript variants. [provided by RefSeq, Jun 2011]






Synonyms: CD56; MSK39; NCAM

**Protein Families:** Druggable Genome, ES Cell Differentiation/IPS, Transmembrane

**Protein Pathways:** Cell adhesion molecules (CAMs), Prion diseases

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NCAM1 ([RC207890], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NCAM1. Positive lysates [LY405745] (100ug) and [LC405745] (20ug) can be purchased separately from OriGene.