

Product datasheet for TA805112S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Nkx3.1 (NKX3-1) Mouse Monoclonal Antibody [Clone ID: OTI4B4]

Product data:

Isotype:

Product Type: Primary Antibodies

Clone Name: OTI4B4
Applications: IHC, WB
Recommended Dilution: WB 1:2000
Reactivity: Human
Host: Mouse

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human NKX3-1 (NP_006158) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

lgG1

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 26.2 kDa

Gene Name: NK3 homeobox 1

Database Link: NP 006158

Entrez Gene 4824 Human

Q99801

Background: This gene encodes a homeobox-containing transcription factor. This transcription factor

functions as a negative regulator of epithelial cell growth in prostate tissue. Aberrant expression of this gene is associated with prostate tumor progression. Alternate splicing

results in multiple transcript variants of this gene. [provided by RefSeq, Jan 2012]

Synonyms: BAPX2; NKX3; NKX3.1; NKX3A


Protein Families: Druggable Genome, Transcription Factors






Protein Pathways: Pathways in cancer, Prostate cancer

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY NKX3-1 ([RC210374], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NKX3-1. Positive lysates [LY401858] (100ug) and [LC401858] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-NKX3-1 monoclonal antibody (1:200).





Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-NKX3-1 mouse monoclonal antibody. ([TA805112]) Dilution: 1:150