

Product datasheet for TA805325AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

ASPA Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1A11]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1A11

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 77-313 of human ASPA

(NP_000040) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 35.6 kDa

Gene Name: aspartoacylase

Database Link: NP 000040

Entrez Gene 11484 MouseEntrez Gene 79251 RatEntrez Gene 443 Human

P45381

Background: This gene encodes an enzyme that catalyzes the conversion of N-acetyl L-aspartic acid (NAA)

to aspartate and acetate. NAA is abundant in the brain where hydrolysis by aspartoacylase is thought to help maintain white matter. This protein is an NAA scavenger in other tissues. Mutations in this gene cause Canavan disease. Alternatively spliced transcript variants have

been found for this gene. [provided by RefSeq, Jul 2008]






Synonyms: ACY2; ASP


Protein Families: Druggable Genome

Protein Pathways: Alanine, aspartate and glutamate metabolism, Histidine metabolism

Product images:



Immunohistochemical staining of paraffinembedded Human liver tissue within the normal limits using anti-ASPA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA805325])



Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-ASPA mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA805325])