

Product datasheet for TA805591M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Caspase 4 (CASP4) Mouse Monoclonal Antibody [Clone ID: OTI9F6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI9F6

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 81-270 of human CASP4

(NP 150649) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 36.5 kDa

Gene Name: caspase 4

Database Link: NP 150649

Entrez Gene 837 Human

P49662




Background:

This gene encodes a protein that is a member of the cysteine-aspartic acid protease (caspase) family. Sequential activation of caspases plays a central role in the execution-phase of cell apoptosis. Caspases exist as inactive proenzymes composed of a prodomain and a large and small protease subunit. Activation of caspases requires proteolytic processing at conserved internal aspartic residues to generate a heterodimeric enzyme consisting of the large and small subunits. This caspase is able to cleave and activate its own precursor protein, as well as caspase 1 precursor. When overexpressed, this gene induces cell apoptosis. Alternative splicing results in transcript variants encoding distinct isoforms. [provided by RefSeq, Jul 2008]

Synonyms: ICE(rel)II; ICEREL-II; ICH-2; Mih1; Mih1/TX; TX

Protein Families: Druggable Genome, Protease

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY CASP4 ([RC204711], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CASP4. Positive lysates [LY409623] (100ug) and [LC409623] (20ug) can be purchased separately from OriGene.