

Product datasheet for TA805653BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Rb (RB1) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI11C5]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI11C5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 309-590 of human

RB1(NP 000312) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: RB transcriptional corepressor 1

Database Link: NP 000312

Entrez Gene 19645 MouseEntrez Gene 24708 RatEntrez Gene 5925 Human

P06400

Background: The protein encoded by this gene is a negative regulator of the cell cycle and was the first

tumor suppressor gene found. The encoded protein also stabilizes constitutive

heterochromatin to maintain the overall chromatin structure. The active,

hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene

are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic

sarcoma. [provided by RefSeq, Jul 2008]

Synonyms: OSRC; p105-Rb; pp110; PPP1R130; pRb; RB






Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: Bladder cancer, Cell cycle, Chronic myeloid leukemia, Glioma, Melanoma, Non-small cell lung

cancer, Pancreatic cancer, Pathways in cancer, Prostate cancer, Small cell lung cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RB1 ([RC206933], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RB1 mouse monoclonal antibody.1:500`