

## Product datasheet for TA805799M

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Interferon gamma (IFNG) Mouse Monoclonal Antibody [Clone ID: OTI4D7]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI4D7

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human Host: Mouse

Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 24-166 of human

IFNG(NP 000610) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 16.7 kDa

**Gene Name:** interferon, gamma

Database Link: NP 000610

Entrez Gene 3458 Human

P01579

**Background:** This gene encodes a member of the type II interferon family. The protein encoded is a

soluble cytokine with antiviral, immunoregulatory and anti-tumor properties and is a potent

activator of macrophages. Mutations in this gene are associated with aplastic anemia.

[provided by RefSeq, Nov 2009]

Synonyms: IFG; IFI






**Protein Families:** Druggable Genome, Secreted Protein

Protein Pathways: Allograft rejection, Cytokine-cytokine receptor interaction, Graft-versus-host disease, Jak-STAT

signaling pathway, Natural killer cell mediated cytotoxicity, Proteasome, Regulation of autophagy, Systemic lupus erythematosus, T cell receptor signaling pathway, TGF-beta

signaling pathway, Type I diabetes mellitus

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IFNG ([RC209993], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IFNG (1:2000). Positive lysates [LY400207] (100ug) and [LC400207] (20ug) can be purchased separately from OriGene.