

## **Product datasheet for TA805843**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# **ROS1 Mouse Monoclonal Antibody [Clone ID: OTI1E6]**

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI1E6
Applications: IHC, WB

Recommended Dilution: WB 1:500-2000, IHC 1:5000

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 2126-2347 of human

ROS1(NP 002935) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Gene Name:** ROS proto-oncogene 1, receptor tyrosine kinase

Database Link: NP 002935

Entrez Gene 6098 Human

P08922

Background: This proto-oncogene, highly-expressed in a variety of tumor cell lines, belongs to the

sevenless subfamily of tyrosine kinase insulin receptor genes. The protein encoded by this gene is a type I integral membrane protein with tyrosine kinase activity. The protein may function as a growth or differentiation factor receptor. [provided by RefSeq, Jul 2008]


**Synonyms:** c-ros-1; MCF3; ROS

**Protein Families:** Druggable Genome, Protein Kinase, Transmembrane







### **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY ROS1 ([RC220652], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ROS1.



Western blot analysis of extracts (35ug) from 2 different cell lines by using anti-ROS1 monoclonal antibody (1:500).



Immunohistochemical staining of paraffinembedded HCC78 (left) and HeLa (right) xenograft using ROS1 (OTI1A1) mAb. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.