

Product datasheet for TA805908AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MTOR Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C3
Applications: IHC, WB
Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1766-2144 of human

MTOR(NP 004949) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 288.7 kDa

Gene Name: mechanistic target of rapamycin

Database Link: NP 004949

Entrez Gene 56717 MouseEntrez Gene 56718 RatEntrez Gene 2475 Human

P42345

Synonyms: FRAP; FRAP1; FRAP2; RAFT1; RAPT1; SKS

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Acute myeloid leukemia, Adipocytokine signaling pathway, ErbB signaling pathway, Glioma,


Insulin signaling pathway, mTOR signaling pathway, Pathways in cancer, Prostate cancer,

Type II diabetes mellitus






Product images:




Human recombinant protein fragment corresponding to amino acids 1766-2144 of human MTOR (NP_004949) produced in E.coli.




Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-MTOR mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA805908]) (1:500)





Immunohistochemical staining of paraffinembedded Human lymphoma tissue using anti-MTOR mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA805908]) (1:500)



Immunohistochemical staining of paraffinembedded Human spleen tissue within the normal limits using anti-MTOR mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA805908]) (1:500)