

Product datasheet for TA806157M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Steroidogenic Factor 1 (NR5A1) Mouse Monoclonal Antibody [Clone ID: OTI1B4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1B4
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse IgG2a

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-262 of human

NR5A1(NP_004950) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 51.5 kDa

Gene Name: nuclear receptor subfamily 5 group A member 1

Database Link: NP 004950

Entrez Gene 26423 MouseEntrez Gene 83826 RatEntrez Gene 2516 Human

O13285


Synonyms: AD4BP; ELP; FTZ1; FTZF1; hSF-1; POF7; SF-1; SFGF8; SRXY3

Protein Families: Druggable Genome, Transcription Factors






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NR5A1 (Cat# [RC207577], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NR5A1(Cat# [TA806157]). Positive lysates [LY417628] (100ug) and [LC417628] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-NR5A1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.





Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-NR5A1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.