

Product datasheet for TA806413M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TrkA (NTRK1) Mouse Monoclonal Antibody [Clone ID: OTI5B6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI5B6
Applications: IHC, WB

Reactivity: WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 440-796 of human

NTRK1(NP_002520) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 87.3 kDa

Gene Name: neurotrophic receptor tyrosine kinase 1

Database Link: NP 002520

Entrez Gene 18211 MouseEntrez Gene 59109 RatEntrez Gene 4914 Human

P04629





Background:

This gene encodes a member of the neurotrophic tyrosine kinase receptor (NTKR) family. This kinase is a membrane-bound receptor that, upon neurotrophin binding, phosphorylates itself and members of the MAPK pathway. The presence of this kinase leads to cell differentiation and may play a role in specifying sensory neuron subtypes. Mutations in this gene have been associated with congenital insensitivity to pain, anhidrosis, self-mutilating behavior, mental retardation and cancer. Alternate transcriptional splice variants of this gene have been found, but only three have been characterized to date. [provided by RefSeq, Jul 2008]


Synonyms: MTC; p140-TrkA; TRK; Trk-A; TRK1; TRKA

Protein Families: Druggable Genome, Protein Kinase, Transmembrane


Protein Pathways: Apoptosis, Endocytosis, MAPK signaling pathway, Neurotrophin signaling pathway, Pathways

in cancer, Thyroid cancer

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY NTRK1 (Cat# [RC213047], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-NTRK1(Cat# [TA806413]). Positive lysates [LY419273] (100ug) and [LC419273] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Carcinoma of Human pancreas tissue using anti-NTRK1 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.