

Product datasheet for TA806450M

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

RHOC Mouse Monoclonal Antibody [Clone ID: OTI3C6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C6

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Synthetic peptide corresponding to C terminal of human RHOC.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 21.8 kDa

Gene Name: ras homolog family member C

Database Link: NP 786886

Entrez Gene 11853 MouseEntrez Gene 295342 RatEntrez Gene 389 Human

P08134






Background:

This gene encodes a member of the Rho family of small GTPases, which cycle between inactive GDP-bound and active GTP-bound states and function as molecular switches in signal transduction cascades. Rho proteins promote reorganization of the actin cytoskeleton and regulate cell shape, attachment, and motility. The protein encoded by this gene is prenylated at its C-terminus, and localizes to the cytoplasm and plasma membrane. It is thought to be important in cell locomotion. Overexpression of this gene is associated with tumor cell proliferation and metastasis. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul 2008]

Synonyms: ARH9; ARHC; H9; RHOH9

Protein Families: Druggable Genome

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY RHOC ([RC217556], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-RHOC (1:2000). Positive lysates [LY403579] (100ug) and [LC403579] (20ug) can be purchased separately from OriGene.