

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# Product datasheet for TA806464S

## Factor XIII (F13B) Mouse Monoclonal Antibody [Clone ID: OTI2D4]

## **Product data:**

| Product Type:           | Primary Antibodies  |
|-------------------------|---|
| Clone Name:             | OTI2D4  |
| Applications:           | WB  |
| Recommended Dilution:   | WB 1:500  |
| Reactivity:             | Human   |
| Host:                   | Mouse   |
| lsotype:                | lgG1  |
| Clonality:              | Monoclonal  |
| Immunogen:              | Human recombinant protein fragment corresponding to amino acids 357-661 of human<br>F13B(NP_001985) produced in E.coli. |
| Formulation:            | PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.  |
| Concentration:          | 1 mg/ml   |
| Purification:           | Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography<br>(protein A/G)            |
| Conjugation:            | Unconjugated  |
| Storage:                | Store at -20°C as received.   |
| Stability:              | Stable for 12 months from date of receipt.  |
| Predicted Protein Size: | 73.2 kDa  |
| Gene Name:              | coagulation factor XIII B chain   |
| Database Link:          | <u>NP_001985</u><br><u>Entrez Gene 2165 Human</u><br><u>P05160</u>  |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

### Sactor XIII (F13B) Mouse Monoclonal Antibody [Clone ID: OTI2D4] – TA806464S

Background:This gene encodes coagulation factor XIII B subunit. Coagulation factor XIII is the last<br/>zymogen to become activated in the blood coagulation cascade. Plasma factor XIII is a<br/>heterotetramer composed of 2 A subunits and 2 B subunits. The A subunits have catalytic<br/>function, and the B subunits do not have enzymatic activity and may serve as a plasma carrier<br/>molecules. Platelet factor XIII is comprised only of 2 A subunits, which are identical to those<br/>of plasma origin. Upon activation by the cleavage of the activation peptide by thrombin and<br/>in the presence of calcium ion, the plasma factor XIII dissociates its B subunits and yields the<br/>same active enzyme, factor XIIIa, as platelet factor XIII. This enzyme acts as a<br/>transglutaminase to catalyze the formation of gamma-glutamyl-epsilon-lysine crosslinking<br/>between fibrin molecules, thus stabilizing the fibrin clot. Factor XIII deficiency is classified into<br/>two categories: type I deficiency, characterized by the lack of both the A and B subunits; and<br/>type II deficiency, characterized by the lack of the A subunit alone. These defects can result in<br/>a lifelong bleeding tendency, defective wound healing, and habitual abortion. [provided by<br/>RefSeq, Jul

| Synonyms:         | FXIIIB                              |
|-------------------|-------------------------------------|
| Protein Families: | Druggable Genome                    |
| Protein Pathways: | Complement and coagulation cascades |

### **Product images:**

| 170 | _ | - |
|-----|---|---|
| 130 | - |   |
| 100 | _ | - |
| 70  | _ |   |
| 55  |   |   |
| 40  |   |   |
| 35  |   |   |
| 25  | - |   |
| 15  | - |   |
| 10  | - |   |
|     |   |   |

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY F13B ([RC219609], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-F13B. Positive lysates [LY419598] (100ug) and [LC419598] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US