

Product datasheet for TA806504M

UBE2D2 Mouse Monoclonal Antibody [Clone ID: OTI1H4]

Product data:

Product Type:	Primary Antibodies	
Clone Name:	OTI1H4	
Applications:	WB	
Recommended Dilution:	WB 1:2000	
Reactivity:	Human, Mouse, Rat	
Host:	Mouse	
lsotype:	lgG2b	
Clonality:	Monoclonal	
Immunogen:	Full length human recombinant protein of human UBE2D2 (NP_003330) produced in E.coli.	
Formulation:	PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.	
Concentration:	1 mg/ml	
Purification:	Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography (protein A/G)	
Conjugation:	Unconjugated	
Storage:	Store at -20°C as received.	
Stability:	Stable for 12 months from date of receipt.	
Predicted Protein Size:	16.6 kDa	
Gene Name:	ubiquitin conjugating enzyme E2 D2	
Database Link:	<u>NP_003330</u> <u>Entrez Gene 641452 RatEntrez Gene 7322 Human</u> <u>P62837</u>	



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

	UBE2D2 Mouse Monoclonal Antibody [Clone ID: OTI1H4] – TA806504M	
Background:	Regulated degradation of misfolded, damaged or short-lived proteins in eukaryotes occurs via the ubiquitin (Ub)-proteasome system (UPS). An integral part of the UPS system is the ubiquitination of target proteins and covalent linkage of Ub-containing proteins to form polymeric chains, marking them as targets for 26S proteasome-mediated degradation. Ubiquitination of proteins is mediated by a cascade of enzymes which includes E1 (ubiquitin activating), E2 (ubiquitin conjugating), and E3 (ubiquitin ligases) enzymes. This gene encodes a member of the E2 enzyme family. Substrates of this enzyme include the tumor suppressor protein p53 and peroxisomal biogenesis factor 5 (PEX5). Alternative splicing results in multiple transcript variants of this gene. [provided by RefSeq, May 2013]	
Synonyms:	E2(17)KB2; PUBC1; UBC4; UBC4/5; UBCH4; UBCH5B	
Protein Pathway	Ubiquitin mediated proteolysis	

Product images:

170	_	
130	_	
100	_	
70		
55		
40		
35	—	
25	—	
15	—	
10	_	

HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY UBE2D2 ([RC218795], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-UBE2D2. Positive lysates [LY418757] (100ug) and [LC418757] (20ug) can be purchased separately from OriGene.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2025 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US