

Product datasheet for TA806648M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

INDOL1 (IDO2) Mouse Monoclonal Antibody [Clone ID: OTI19F8]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI19F8
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human IDO2 (NP_919270) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Gene Name: indoleamine 2,3-dioxygenase 2

Database Link: NP 919270

Entrez Gene 169355 Human

Q6ZQW0

Background: Along with the enzymes encoded by the INDO (MIM 147435) and TDO2 (MIM 191070) genes,

the enzyme encoded by the INDOL1 gene metabolizes tryptophan in the kynurenine pathway

(Ball et al., 2007 [PubMed 17499941]). [supplied by OMIM, Feb 2011]


Synonyms: INDOL1

Protein Pathways: Metabolic pathways, Tryptophan metabolism






Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY IDO2 ([RC223337], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-IDO2. Positive lysates [LY403659] (100ug) and [LC403659] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human tonsil within the normal limits using anti-IDO2 mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.