

# **Product datasheet for TA806794**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

# JAK2 Mouse Monoclonal Antibody [Clone ID: OTI7D12]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI7D12

Applications: IHC

Recommended Dilution: IHC 1:150

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 297-596 of human

JAK2(NP\_004963) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Gene Name: Janus kinase 2

Database Link: NP 004963

Entrez Gene 16452 MouseEntrez Gene 24514 RatEntrez Gene 3717 Human

<u>060674</u>

**Background:** This gene product is a protein tyrosine kinase involved in a specific subset of cytokine

receptor signaling pathways. It has been found to be constituitively associated with the prolactin receptor and is required for responses to gamma interferon. Mice that do not express an active protein for this gene exhibit embryonic lethality associated with the

absence of definitive erythropoiesis. [provided by RefSeq, Jul 2008]

Synonyms: JTK10; THCYT3







**Protein Families:** Druggable Genome, Protein Kinase

**Protein Pathways:** Adipocytokine signaling pathway, Chemokine signaling pathway, Jak-STAT signaling pathway

# **Product images:**



Immunohistochemical staining of paraffinembedded Human prostate tissue within the normal limits using anti-JAK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA806794) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-JAK2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA806794) (1:150)