

Product datasheet for TA806835S

OriGene Technologies, Inc.9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: OTI8B1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI8B1

Applications: ELISA, FC, IF, LMNX, Neutralize, WB

Recommended Dilution: WB 1:2000, IF 1:100~900, FLOW 1:50

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDCD1 (NP_005009) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29.2 kDa

Gene Name: programmed cell death 1

Database Link: NP 005009

Entrez Gene 5133 Human

Q15116


Synonyms: CD279; hPD-1; hPD-l; hSLE1; PD-1; PD1; SLEB2

Protein Families: Druggable Genome, Transmembrane


Protein Pathways: Cell adhesion molecules (CAMs), T cell receptor signaling pathway




Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1 ([RC210364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1. Positive lysates [LY401555] (100ug) and [LC401555] (20ug) can be purchased separately from OriGene.




PD1 ELISA with 1B11 Capture ([CF807867]) and 8B1 Detection ([CF806835]) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])





Immunofluorescent staining of PDCD1 ([RC210364])-stable-transfected HEK293T cells (left) labeling PDCD1 with mouse monoclonal antibody [TA806835] (green) and nucleus with Hoechst33342 (blue). HEK293T cells serve as negative control (right). n (1:100)






293T cell line






Immunocytochemistry staining of stable expression PD1 cells using anti-PDCD1 mouse monoclonal antibody ([TA806835]) (Left). The rihgt is negative control. (1:100) (1:900)




Flow cytometric Analysis of stable expression PD1 cells using anti-PDCD1 antibody ([TA806835]) (Red) compared to a nonspecific negative control antibody (Blue) (1:50).




HEK293T cells transfected with either mouse PD1 ([MR227347]) overexpress plasmid (Red) or empty vector control plasmid (Blue) were immunostained by anti-PDCD1 antibody ([TA806835]), compared to a nonspecific negative control antibody (green), and then analyzed by flow cytometry (1:50).






PD1 Luminex ELISA with 1B11 Capture ([CF807867]) and 8B1 Detection ([CF806835]) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])



PD1 Luminex ELISA with 7B4 Capture ([CF807995]) and 8B1 Detection ([CF806835]) Antibodies. Substrate used: Recombinant Human PD1 ([TP310364])



Flow cytometric Analysis of stable expression PDL1 ([RC213071]) cells using anti-PDCD1 antibody ([TA806835]) (blue) or 0.3ug/ml PD1-Fc fusion protein ([TP700199]) (green) or both (red), and detected by anti-Fc (human) IgG-FITC (1:50).