

Product datasheet for TA806858AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

VIP Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI1C2]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1C2
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human VIP (NP_003372) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 16.4 kDa

Gene Name: vasoactive intestinal peptide

Database Link: NP 003372

Entrez Gene 7432 Human

P01282

Background: The protein encoded by this gene belongs to the glucagon family. It stimulates myocardial

contractility, causes vasodilation, increases glycogenolysis, lowers arterial blood pressure and relaxes the smooth muscle of trachea, stomach and gall bladder. Alternative splicing occurs at

this locus and two transcript variants encoding distinct isoforms have been identified.

[provided by RefSeq, Jul 2008]

Synonyms: PHM27







Protein Families:

Druggable Genome, Secreted Protein, Transmembrane

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY VIP ([RC203126], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-VIP. Positive lysates [LY418679] (100ug) and [LC418679] (20ug) can be purchased separately from OriGene.




Immunohistochemical staining of paraffinembedded Human liver tissue using anti-VIP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA806858]) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-VIP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA806858]) (1:150)



Immunohistochemical staining of paraffinembedded Carcinoma of Human prostate tissue using anti-VIP mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, [TA806858]) (1:150)