

Product datasheet for TA806866M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

c Fos (FOS) Mouse Monoclonal Antibody [Clone ID: OTI4A10]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI4A10

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human FOS (NP_005243) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 40.5 kDa

Gene Name: Fos proto-oncogene, AP-1 transcription factor subunit

Database Link: NP 005243

Entrez Gene 14281 MouseEntrez Gene 314322 RatEntrez Gene 2353 Human

P01100

Background: The Fos gene family consists of 4 members: FOS, FOSB, FOSL1, and FOSL2. These genes

encode leucine zipper proteins that can dimerize with proteins of the JUN family, thereby forming the transcription factor complex AP-1. As such, the FOS proteins have been implicated as regulators of cell proliferation, differentiation, and transformation. In some cases, expression of the FOS gene has also been associated with apoptotic cell death.

[provided by RefSeg, Jul 2008]






Synonyms: AP-1; C-FOS; p55

Protein Families: Druggable Genome, Transcription Factors

Protein Pathways: B cell receptor signaling pathway, Colorectal cancer, MAPK signaling pathway, Pathways in

cancer, T cell receptor signaling pathway, Toll-like receptor signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY FOS ([RC202597], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-FOS. Positive lysates [LY401614] (100ug) and [LC401614] (20ug) can be purchased separately from OriGene.