

Product datasheet for TA807065

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

TEM1 (CD248) Mouse Monoclonal Antibody [Clone ID: OTI1H6]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI1H6
Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 20-300 of human

CD248(NP_065137) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 80.7 kDa

Gene Name: CD248 molecule

Database Link: NP 065137

Entrez Gene 57124 Human

Q9HCU0


Synonyms: CD164L1; TEM1

Protein Families: Druggable Genome, Transmembrane






Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CD248 (Cat# [RC208576], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CD248 (Cat# TA807065)(1:500). Positive lysates [LY412436] (100ug) and [LC412436] (20ug) can be purchased separately from OriGene.



Immunohistochemical staining of paraffinembedded Human Kidney tissue within the normal limits using anti-CD248 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807065) (1:150)





Immunohistochemical staining of paraffinembedded Carcinoma of Human liver tissue using anti-CD248 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807065) (1:150)