

Product datasheet for TA807075M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

CITED1 Mouse Monoclonal Antibody [Clone ID: OTI3C12]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C12
Applications: IHC, WB

Recommended Dilution: WB 1:2000, IHC 1:150

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human CITED1 (NP_004134) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 19.7 kDa

Gene Name: Cbp/p300 interacting transactivator with Glu/Asp rich carboxy-terminal domain 1

Database Link: NP 004134

Entrez Gene 4435 Human

Q99966

Background: This gene encodes a member of the CREB-binding protein/p300-interacting transactivator

with Asp/Glu-rich C-terminal domain (CITED) family of proteins. The encoded protein, also known as melanocyte-specific gene 1, may function as a transcriptional coactivator and may play a role in pigmentation of melanocytes. Alternatively spliced transcript variants have

been described. [provided by RefSeq, Jan 2009]


Synonyms: MSG1






Protein Families: Transcription Factors

Product images:




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY CITED1 (Cat# [RC202419], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-CITED1(Cat# [TA807075]). Positive lysates [LY418187] (100ug) and [LC418187] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (35ug) from 9 different cell lines by using anti-CITED1 monoclonal antibody (HepG2: human; HeLa: human; SVT2: mouse; A549: human; COS7: monkey; Jurkat: human; MDCK: canine; PC12: rat; MCF7: human) (1:200).





Immunohistochemical staining of paraffinembedded Human testicular cancer tissue using anti-CITED1 mouse monoclonal antibody. Heatinduced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.