

Product datasheet for TA807098

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436

OriGene Technologies, Inc.

Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PD1 (PDCD1) Mouse Monoclonal Antibody [Clone ID: OTI26C4]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI26C4

Applications: FC, IF, Neutralize, WB

Recommended Dilution: WB 1:2000, IF 1:100~900, FLOW 1:50

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human PDCD1 (NP_005009) produced in HEK293T

cell

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 29.2 kDa

Gene Name: programmed cell death 1

Database Link: NP 005009

Entrez Gene 5133 Human

015116



Background:


This gene encodes a cell surface membrane protein of the immunoglobulin superfamily. This protein is expressed in pro-B-cells and is thought to play a role in their differentiation. In mice, expression of this gene is induced in the thymus when anti-CD3 antibodies are injected and large numbers of thymocytes undergo apoptosis. Mice deficient for this gene bred on a BALB/c background developed dilated cardiomyopathy and died from congestive heart failure. These studies suggest that this gene product may also be important in T cell function and contribute to the prevention of autoimmune diseases. [provided by RefSeq, Jul 2008]

Synonyms: CD279; hPD-1; hPD-1; hSLE1; PD-1; PD1; SLEB2


Protein Families: Druggable Genome, Transmembrane

Protein Pathways: Cell adhesion molecules (CAMs), T cell receptor signaling pathway

Product images:





HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PDCD1 ([RC210364], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PDCD1 (1:2000). Positive lysates [LY401555] (100ug) and [LC401555] (20ug) can be purchased separately from OriGene.




Immunofluorescent staining of PDCD1 ([RC210364])-stable-transfected HEK293T cells (left) labeling PDCD1 with mouse monoclonal antibody TA807098 (green) and nucleus with Hoechst33342 (blue). HEK293T cells serve as negative control (right) (1:100).











Immunocytochemistry staining of stable expression PD1 cells using anti-PDCD1 mouse monoclonal antibody (TA807098) (Left). The rihgt is negative control. (1:100) (1:900)



Flow cytometric Analysis of stable expression PD1 cells using anti-PDCD1 antibody (TA807098) (Red) compared to a nonspecific negative control antibody (Blue) (1:50).



Flow cytometric Analysis of stable expression PDL1 ([RC213071]) cells using anti-PDCD1 antibody (TA807098) (blue) or 0.3ug/ml PD1-Fc fusion protein ([TP700199]) (green) or both (red), and detected by anti-Fc (human) IgG-FITC (1:50).