

# **Product datasheet for TA807106**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Renin (REN) Mouse Monoclonal Antibody [Clone ID: OTI2A2]

### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2A2

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human
Host: Mouse
Isotype: IgG2b

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 131-406 of human

REN(NP 000528) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 42.3 kDa **Gene Name:** renin

Database Link: NP 000528

Entrez Gene 5972 Human

P00797



Background:


Renin catalyzes the first step in the activation pathway of angiotensinogen--a cascade that can result in aldosterone release, vasoconstriction, and increase in blood pressure. Renin, an aspartyl protease, cleaves angiotensinogen to form angiotensin I, which is converted to angiotensin II by angiotensin I converting enzyme, an important regulator of blood pressure and electrolyte balance. Transcript variants that encode different protein isoforms and that arise from alternative splicing and the use of alternative promoters have been described, but their full-length nature has not been determined. Mutations in this gene have been shown to cause familial hyperproreninemia. [provided by RefSeq, Jul 2008]

Synonyms: ADTKD4; HNFJ2; RTD

**Protein Families:** Druggable Genome, Secreted Protein

**Protein Pathways:** Renin-angiotensin system

# **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY REN ([RC208382], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-REN (1:2000). Positive lysates [LY424660] (100ug) and [LC424660] (20ug) can be purchased separately from OriGene.