

Product datasheet for TA807167M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

APOB48R (APOBR) Mouse Monoclonal Antibody [Clone ID: OTI6D3]

Product data:

Applications:

Product Type: Primary Antibodies

Clone Name: **OTI6D3** IHC

Recommended Dilution: IHC 1:150

Reactivity: Human Host: Mouse Isotype: lgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 780-1088 of human

APOBR(NP 061160) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Store at -20°C as received. Storage:

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 114.7 kDa

Gene Name: apolipoprotein B receptor

Database Link: NP 061160

Entrez Gene 55911 Human

O0VD83

Background: Apolipoprotein B48 receptor is a macrophage receptor that binds to the apolipoprotein B48

> of dietary triglyceride (TG)-rich lipoproteins. This receptor may provide essential lipids, lipidsoluble vitamins and other nutrients to reticuloendothelial cells. If overwhelmed with elevated plasma triglyceride, the apolipoprotein B48 receptor may contribute to foam cell formation,

endothelial dysfunction, and atherothrombogenesis. [provided by RefSeq, Jul 2008]







Synonyms: APOB48R; APOB100R

Protein Families: Druggable Genome

Product images:



Immunohistochemical staining of paraffinembedded Human colon tissue within the normal limits using anti-APOBR mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.



Immunohistochemical staining of paraffinembedded Human lymph node tissue within the normal limits using anti-APOBR mouse monoclonal antibody. Heat-induced epitope retrieval by EDTA solution buffer pH 8.0 at 120°C for 3 min.