

Product datasheet for TA807357

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Granulin (GRN) Mouse Monoclonal Antibody [Clone ID: OTI2D1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI2D1

Applications: IHC, WB

Recommended Dilution: WB 1:500, IHC 1:500

Reactivity: Human
Host: Mouse
Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 327-593 of human

GRN(NP_002078) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Unconjugated

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 63.54 kDa

Gene Name: granulin

Database Link: NP 002078

Entrez Gene 2896 Human

P28799


Synonyms: CLN11; GEP; GP88; PCDGF; PEPI; PGRN

Protein Families: Druggable Genome, Secreted Protein






Product images:




Western blot analysis of extracts (35ug) from 4 different cell lines by using anti-GRN monoclonal antibody (1:500).



Equivalent amounts of cell lysates (10 ug per lane) of wild-type 293T cells (WT, Cat# LC810293T) and GRN-Knockout 293T cells (KO, Cat# [LC811733]) were separated by SDS-PAGE and immunoblotted with anti-GRN monoclonal antibody TA807357, (1:100). Then the blotted membrane was stripped and reprobed with antib-actin antibody ([TA811000]) as a loading control.






Immunohistochemical staining of paraffinembedded Human pancreas tissue within the normal limits using anti-GRN mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807357) (1:500)




Immunohistochemical staining of paraffinembedded Human endometrium tissue within the normal limits using anti-GRN mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris buffer (pH8.5) at 120°C for 3min, TA807357) (1:500)



Immunohistochemical staining of paraffinembedded Human spleen tissue within the normal limits using anti-GRN mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA (1:500)





Immunohistochemical staining of paraffinembedded Human appendix tissue within the normal limits using anti-GRN mouse monoclonal antibody. (Heat-induced epitope retrieval by Tris-EDTA (1:0) (1:500)