

Product datasheet for TA807368AM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

PAK4 Mouse Monoclonal Antibody (Biotin conjugated) [Clone ID: OTI3C3]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI3C3

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Human recombinant protein fragment corresponding to amino acids 1-308 of human

PAK4(NP_005875) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 63.9 kDa

Gene Name: p21 (RAC1) activated kinase 4

Database Link: NP 005875

Entrez Gene 70584 MouseEntrez Gene 292756 RatEntrez Gene 10298 Human

096013





Background: PAK proteins, a family of serine/threonine p21-activating kinases, include PAK1, PAK2, PAK3

and PAK4. PAK proteins are critical effectors that link Rho GTPases to cytoskeleton

reorganization and nuclear signaling. They serve as targets for the small GTP binding proteins Cdc42 and Rac and have been implicated in a wide range of biological activities. PAK4

interacts specifically with the GTP-bound form of Cdc42Hs and weakly activates the JNK family

of MAP kinases. PAK4 is a mediator of filopodia formation and may play a role in the reorganization of the actin cytoskeleton. Multiple alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

effector for Cdc42Hs; p21 protein (Cdc42; p21(CDKN1A)-activated kinase 4; p21-activated Synonyms:


kinase 4; protein kinase related to S. cerevisiae STE20; Rac)-activated kinase 4

Protein Families: Druggable Genome, Protein Kinase

Protein Pathways: Axon guidance, ErbB signaling pathway, Focal adhesion, Regulation of actin cytoskeleton,

Renal cell carcinoma, T cell receptor signaling pathway

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PAK4 ([RC202302], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PAK4. Positive lysates [LY417002] (100ug) and [LC417002] (20ug) can be purchased separately from OriGene.