

### **Product datasheet for TA807408S**

# **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

## ECT2 Mouse Monoclonal Antibody [Clone ID: OTI2D9]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI2D9
Applications: IHC, WB

**Reactivity:** WB 1:2000, IHC 1:150 Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 175-438 of human

ECT2(NP\_060568) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol and 0.02% sodium azide.

Concentration: 1 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

**Conjugation:** Unconjugated

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 99.9 kDa

**Gene Name:** epithelial cell transforming 2

Database Link: NP 060568

Entrez Gene 13605 MouseEntrez Gene 361921 RatEntrez Gene 1894 Human

O9H8V3






#### Background:

The protein encoded by this gene is a guanine nucleotide exchange factor and transforming protein that is related to Rho-specific exchange factors and yeast cell cycle regulators. The expression of this gene is elevated with the onset of DNA synthesis and remains elevated during G2 and M phases. In situ hybridization analysis showed that expression is at a high level in cells undergoing mitosis in regenerating liver. Thus, this protein is expressed in a cell cycle-dependent manner during liver regeneration, and is thought to have an important role in the regulation of cytokinesis. Several transcript variants encoding two different isoforms have been found for this gene. [provided by RefSeq, Apr 2012]


Synonyms: ARHGEF31

**Protein Families:** Druggable Genome

## **Product images:**




HEK293T cells were transfected with the pCMV6-ENTRY control (Cat# [PS100001], Left lane) or pCMV6-ENTRY ECT2 (Cat# [RC211319], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-ECT2(Cat# [TA807408]). Positive lysates [LY413314] (100ug) and [LC413314] (20ug) can be purchased separately from OriGene.



Western blot analysis of extracts (10ug) from 1 cell line by using anti-ECT2 monoclonal antibody (1:200).





Immunohistochemical staining of paraffinembedded Adenocarcinoma of Human ovary tissue using anti-ECT2 mouse monoclonal antibody. (Heat-induced epitope retrieval by 1mM EDTA in 10mM Tris, pH8.5, 120°C for 3min, [TA807408])