

# **Product datasheet for TA807428BM**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## PON3 Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI5B5]

#### **Product data:**

**Product Type:** Primary Antibodies

Clone Name: OTI5B5

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

**Immunogen:** Human recombinant protein fragment corresponding to amino acids 30-319 of human

PON3(NP 000931) produced in E.coli.

**Formulation:** PBS (pH 7.3) containing 1% BSA, 50% glycerol.

**Concentration:** 0.5 mg/ml

**Purification:** Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

**Storage:** Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

**Predicted Protein Size:** 39.4 kDa

**Gene Name:** paraoxonase 3

Database Link: NP 000931

Entrez Gene 5446 Human

015166





Background:


This gene is a member of the paraoxonase family and lies in a cluster on chromosome 7 with the other two family members. The encoded protein is secreted into the bloodstream and associates with high-density lipoprotein (HDL). The protein also rapidly hydrolyzes lactones and can inhibit the oxidation of low-density lipoprotein (LDL), a function that is believed to slow the initiation and progression of atherosclerosis. Alternatively spliced variants which encode different protein isoforms have been described; however, only one has been fully characterized. [provided by RefSeq, Jul 2008]

**Synonyms:** lactonase 3; paraoxanase-3; paraoxonase 3; serum paraoxonase

**Protein Families:** Secreted Protein, Transmembrane

**Protein Pathways:** Metabolic pathways

## **Product images:**



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY PON3 ([RC210019], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-PON3. Positive lysates [LY400339] (100ug) and [LC400339] (20ug) can be purchased separately from OriGene.