

Product datasheet for TA807445BM

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

KChIP2 (KCNIP2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI10H1]

Product data:

Product Type: Primary Antibodies

Clone Name: OTI10H1

Applications: WB

Recommended Dilution: WB 1:2000

Reactivity: Human, Mouse, Rat

Host: Mouse Isotype: IgG1

Clonality: Monoclonal

Immunogen: Full length human recombinant protein of human KCNIP2 (NP_775284) produced in E.coli.

Formulation: PBS (pH 7.3) containing 1% BSA, 50% glycerol.

Concentration: 0.5 mg/ml

Purification: Purified from mouse ascites fluids or tissue culture supernatant by affinity chromatography

(protein A/G)

Conjugation: HRP

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Predicted Protein Size: 28.8 kDa

Gene Name: potassium voltage-gated channel interacting protein 2

Database Link: NP 775284

Entrez Gene 56817 RatEntrez Gene 80906 MouseEntrez Gene 30819 Human

Q9NS61





KChIP2 (KCNIP2) Mouse Monoclonal Antibody (HRP conjugated) [Clone ID: OTI10H1] – TA807445BM


Background:

This gene encodes a member of the family of voltage-gated potassium (Kv) channel-interacting proteins (KCNIPs), which belongs to the recoverin branch of the EF-hand superfamily. Members of the KCNIP family are small calcium binding proteins. They all have EF-hand-like domains, and differ from each other in the N-terminus. They are integral subunit components of native Kv4 channel complexes. They may regulate A-type currents, and hence neuronal excitability, in response to changes in intracellular calcium. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified from this gene. [provided by RefSeq, Jul 2008]

Synonyms: KCHIP2

Protein Families: Druggable Genome, Ion Channels: Other

Product images:



HEK293T cells were transfected with the pCMV6-ENTRY control (Left lane) or pCMV6-ENTRY KCNIP2 ([RC203823], Right lane) cDNA for 48 hrs and lysed. Equivalent amounts of cell lysates (5 ug per lane) were separated by SDS-PAGE and immunoblotted with anti-KCNIP2. Positive lysates [LY406638] (100ug) and [LC406638] (20ug) can be purchased separately from OriGene.